

CS675: Convex and Combinatorial Optimization
Fall 2014
Introduction to Matroid Theory

Instructor: Shaddin Dughmi

Optimization over Sets

- Most combinatorial optimization problems can be thought of as choosing the best set from a family of allowable sets
 - Shortest paths
 - Max-weight matching
 - TSP
 - ...

Optimization over Sets

- Most combinatorial optimization problems can be thought of as choosing the best set from a family of allowable sets
 - Shortest paths
 - Max-weight matching
 - TSP
 - ...
- Set system: Pair $(\mathcal{X}, \mathcal{I})$ where \mathcal{X} is a finite **ground set** and $\mathcal{I} \subseteq 2^{\mathcal{X}}$ are the feasible sets

Optimization over Sets

- Most combinatorial optimization problems can be thought of as choosing the best set from a family of allowable sets
 - Shortest paths
 - Max-weight matching
 - TSP
 - ...
- Set system: Pair $(\mathcal{X}, \mathcal{I})$ where \mathcal{X} is a finite **ground set** and $\mathcal{I} \subseteq 2^{\mathcal{X}}$ are the feasible sets
- Objective: often “linear”, referred to as **modular**

Optimization over Sets

- Most combinatorial optimization problems can be thought of as choosing the best set from a family of allowable sets
 - Shortest paths
 - Max-weight matching
 - TSP
 - ...
- Set system: Pair $(\mathcal{X}, \mathcal{I})$ where \mathcal{X} is a finite **ground set** and $\mathcal{I} \subseteq 2^{\mathcal{X}}$ are the feasible sets
- Objective: often “linear”, referred to as **modular**
- Analogues of concave convex: **submodular** and **supermodular** (in no particular order!)

Optimization over Sets

- Most combinatorial optimization problems can be thought of as choosing the best set from a family of allowable sets
 - Shortest paths
 - Max-weight matching
 - TSP
 - ...
- Set system: Pair $(\mathcal{X}, \mathcal{I})$ where \mathcal{X} is a finite **ground set** and $\mathcal{I} \subseteq 2^{\mathcal{X}}$ are the feasible sets
- Objective: often “linear”, referred to as **modular**
- Analogues of concave convex: **submodular** and **supermodular** (in no particular order!)
- Today, we will look only at optimizing modular objectives over an extremely prolific family of set systems
 - Related, directly or indirectly, to a large fraction of optimization problems in P
 - Also pops up in submodular/supermodular optimization problems

Outline


1 Matroids and The Greedy Algorithm

2 Basic Terminology and Properties

3 The Matroid Polytope

4 Matroid Intersection

Maximum Weight Forest Problem



Given a connected undirected graph $G = (V, E)$, and weights $w_e \in \mathbb{R}$ on edges e , find a maximum weight acyclic subgraph (aka **forest**) of G .

- Slight generalization of **minimum weight spanning tree**
- We use n and m to denote $|V|$ and $|E|$, respectively.

The Greedy Algorithm


- 1 $B \leftarrow \emptyset$
- 2 Sort non-negative weight edges in decreasing order of weight
 - e_1, \dots, e_m , with $w_1 \geq w_2 \geq \dots \geq w_m \geq 0$
- 3 For $i = 1$ to m :
 - if $B \cup \{e_i\}$ is acyclic, add e_i to B .

The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
- 2 Sort non-negative weight edges in decreasing order of weight
 - e_1, \dots, e_m , with $w_1 \geq w_2 \geq \dots \geq w_m \geq 0$
- 3 For $i = 1$ to m :
 - if $B \cup \{e_i\}$ is acyclic, add e_i to B .


Theorem

The greedy algorithm outputs a maximum-weight forest.



Lemma


- 1 The empty set is acyclic
- 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
- 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic



Lemma


- 1 The empty set is acyclic
- 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
- 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic

(1) and (2) are trivial, so let's prove (3)




Lemma

- 1 The empty set is acyclic
 - 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
 - 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic
- Sub-lemma: if C is acyclic, then $|C| = n - \#components(C)$.
 - Induction



Lemma


- 1 The empty set is acyclic
 - 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
 - 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic
- Sub-lemma: if C is acyclic, then $|C| = n - \#components(C)$.
 - Induction
 - When $|B| > |A|$, this means $\#components(B) < \#components(A)$



Lemma

- 1 The empty set is acyclic
- 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
- 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic


- Sub-lemma: if C is acyclic, then $|C| = n - \#components(C)$.
 - Induction
- When $|B| > |A|$, this means $\#components(B) < \#components(A)$
- Can't be that all $e \in B$ are "inside" connected components of A



Lemma

- 1 The empty set is acyclic
- 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
- 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic

- Sub-lemma: if C is acyclic, then $|C| = n - \#components(C)$.
 - Induction
- When $|B| > |A|$, this means $\#components(B) < \#components(A)$
- Can't be that all $e \in B$ are "inside" connected components of A
- Some $e \in B$ must "go between" connected components of A .



Lemma

- 1 The empty set is acyclic
 - 2 If A is an acyclic set of edges, and $B \subseteq A$, then B is also acyclic.
 - Converse: if B cyclic then so is A
 - 3 If A, B are acyclic, and $|B| > |A|$, then there is $e \in B \setminus A$ such that $A \cup \{e\}$ is acyclic
 - Inductively: can extend A by adding $|B| - |A|$ elements from $B \setminus A$
- Sub-lemma: if C is acyclic, then $|C| = n - \#components(C)$.
 - Induction
 - When $|B| > |A|$, this means $\#components(B) < \#components(A)$
 - Can't be that all $e \in B$ are "inside" connected components of A
 - Some $e \in B$ must "go between" connected components of A .

Going back to proving the algorithm correct.

Inductive Hypothesis (i)

There is a maximum-weight acyclic forest B_i^* which “agrees” with the algorithm’s choices on edges e_1, \dots, e_i .

- i.e. if B_i denotes the algorithm’s choice up to iteration i , then
$$B_i = B_i^* \cap \{e_1, \dots, e_i\}$$

Going back to proving the algorithm correct.

Inductive Hypothesis (i)

There is a maximum-weight acyclic forest B_i^* which “agrees” with the algorithm’s choices on edges e_1, \dots, e_i .

- i.e. if B_i denotes the algorithm’s choice up to iteration i , then
$$B_i = B_i^* \cap \{e_1, \dots, e_i\}$$

- Assume true for step $i - 1$, and consider step i

Going back to proving the algorithm correct.

Inductive Hypothesis (i)

There is a maximum-weight acyclic forest B_i^* which “agrees” with the algorithm’s choices on edges e_1, \dots, e_i .

- i.e. if B_i denotes the algorithm’s choice up to iteration i , then $B_i = B_i^* \cap \{e_1, \dots, e_i\}$

- Assume true for step $i - 1$, and consider step i
- If $e_i \notin B_i$, then $B_{i-1} \cup \{e_i\}$ is cyclic. Since $B_{i-1} \subseteq B_{i-1}^*$, then $e_i \notin B_{i-1}^*$ (Property 2). So take $B_i^* = B_{i-1}^*$.

Going back to proving the algorithm correct.

Inductive Hypothesis (i)

There is a maximum-weight acyclic forest B_i^* which “agrees” with the algorithm’s choices on edges e_1, \dots, e_i .

- i.e. if B_i denotes the algorithm’s choice up to iteration i , then
$$B_i = B_i^* \cap \{e_1, \dots, e_i\}$$

- Assume true for step $i - 1$, and consider step i
- If $e_i \notin B_i$, then $B_{i-1} \cup \{e_i\}$ is cyclic. Since $B_{i-1} \subseteq B_{i-1}^*$, then $e_i \notin B_{i-1}^*$ (Property 2). So take $B_i^* = B_{i-1}^*$.
- If $e_i \in B_i$ and $e_i \notin B_i^*$, extend B_i to the size of B_{i-1}^* (property 3)
 - Recall that $B_{i-1} = B_i \setminus \{e_i\} \subseteq B_{i-1}^*$
 - $B_i^* = B_{i-1}^* \cup \{e_i\} \setminus \{e_k\}$ for some $k > i$
 - B_i^* has weight no less than B_{i-1}^* , so optimal.

To prove optimality of the greedy algorithm, all we needed was the following.

Matroids

A set system $M = (\mathcal{X}, \mathcal{I})$ is a **matroid** if

- 1 $\emptyset \in \mathcal{I}$
- 2 If $A \in \mathcal{I}$ and $B \subseteq A$, then $B \in \mathcal{I}$ (Downward Closure)
- 3 If $A, B \in \mathcal{I}$ and $|B| > |A|$, then $\exists x \in B \setminus A$ such that $A \cup \{x\} \in \mathcal{I}$ (Exchange Property)

To prove optimality of the greedy algorithm, all we needed was the following.

Matroids

A set system $M = (\mathcal{X}, \mathcal{I})$ is a **matroid** if

- 1 $\emptyset \in \mathcal{I}$
- 2 If $A \in \mathcal{I}$ and $B \subseteq A$, then $B \in \mathcal{I}$ (Downward Closure)
- 3 If $A, B \in \mathcal{I}$ and $|B| > |A|$, then $\exists x \in B \setminus A$ such that $A \cup \{x\} \in \mathcal{I}$ (Exchange Property)

- Three conditions above are called the **matroid axioms**
- $A \in \mathcal{I}$ is called an **independent set** of the matroid.

To prove optimality of the greedy algorithm, all we needed was the following.

Matroids

A set system $M = (\mathcal{X}, \mathcal{I})$ is a **matroid** if

- 1 $\emptyset \in \mathcal{I}$
- 2 If $A \in \mathcal{I}$ and $B \subseteq A$, then $B \in \mathcal{I}$ (Downward Closure)
- 3 If $A, B \in \mathcal{I}$ and $|B| > |A|$, then $\exists x \in B \setminus A$ such that $A \cup \{x\} \in \mathcal{I}$ (Exchange Property)

- Three conditions above are called the **matroid axioms**
- $A \in \mathcal{I}$ is called an **independent set** of the matroid.
- The matroid whose independent sets are acyclic subgraphs is called a **graphic matroid**
- Other examples abound!

Example: Linear Matroid

- \mathcal{X} is a finite set of vectors $\{v_1, \dots, v_m\} \subseteq \mathbb{R}^n$
- $S \in \mathcal{I}$ iff the vectors in S are linearly independent
- Downward closure: If a set of vectors is linearly independent, then every subset of it is also
- Exchange property: Can always extend a low-dimension independent set S by adding vectors from a higher dimension independent set T


Example: Uniform Matroid

- \mathcal{X} is an arbitrary finite set $\{1, \dots, n\}$.
- $S \in \mathcal{I}$ iff $|S| \leq k$.

- Downward closure: If a set S has $|S| \leq k$ then the same holds for $T \subseteq S$.
- Exchange property: If $|S| < |T| \leq k$, then there is an element in $T \setminus S$, and we can add it to S while preserving independence.

Example: Partition Matroid

- \mathcal{X} is the disjoint union of classes X_1, \dots, X_m
 - Each class X_j has an upperbound k_j .
 - $S \in \mathcal{I}$ iff $|S \cap X_j| \leq k_j$ for all j
-
- This is the “disjoint union” of a number of uniform matroids



Example: Transversal Matroid

- Described by a bipartite graph $E \subseteq L \times R$
 - $\mathcal{X} = L$
 - $S \in \mathcal{I}$ iff there is a bipartite matching which matches S
- Downward closure: If we can match S , then we can match $T \subseteq S$.
 - Exchange property: If $|T| > |S|$ is matchable, then an **augmenting path/alternating path** amends the extends the matching of S to some $x \in T \setminus S$.

The Greedy Algorithm on Matroids

The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
- 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
- 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .

Theorem

The greedy algorithm returns the maximum weight set for every choice of weights if and only if the set system $(\mathcal{X}, \mathcal{I})$ is a matroid.

We already saw the “if” direction. We will skip “only if”.

The Greedy Algorithm on Matroids

The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
- 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
- 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .

The Greedy Algorithm on Matroids

The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
 - 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
 - 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .
- To implement this, we need an **independence oracle** for step 3
 - A subroutine which checks whether $S \in \mathcal{I}$ or not.
 - Runs in time $O(n \log n) + nT$, where T is runtime of the independence oracle.

The Greedy Algorithm on Matroids

The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
 - 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
 - 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .
- To implement this, we need an **independence oracle** for step 3
 - A subroutine which checks whether $S \in \mathcal{I}$ or not.
 - Runs in time $O(n \log n) + nT$, where T is runtime of the independence oracle.
 - For most “natural” matroids, independence oracle is easy to implement efficiently
 - Graphic matroid
 - Linear matroid
 - Uniform/partition matroid
 - Transversal matroid

Outline

- 1 Matroids and The Greedy Algorithm
- 2 Basic Terminology and Properties**
- 3 The Matroid Polytope
- 4 Matroid Intersection

Independent Sets, Bases, and Circuits

Consider a matroid $\mathcal{M} = (\mathcal{X}, \mathcal{I})$.

- An **independent set** is a set $A \in \mathcal{I}$.

Independent Sets, Bases, and Circuits

Consider a matroid $\mathcal{M} = (\mathcal{X}, \mathcal{I})$.

- An **independent set** is a set $A \in \mathcal{I}$.
- A **base** of \mathcal{M} is a maximal independent set

Independent Sets, Bases, and Circuits

Consider a matroid $\mathcal{M} = (\mathcal{X}, \mathcal{I})$.

- An **independent set** is a set $A \in \mathcal{I}$.
- A **base** of \mathcal{M} is a maximal independent set
- A **base of** $S \subseteq \mathcal{X}$ in \mathcal{M} is maximal independent subset of S
 - I.e. a base of the matroid after deleting \overline{S} .

Independent Sets, Bases, and Circuits

Consider a matroid $\mathcal{M} = (\mathcal{X}, \mathcal{I})$.

- An **independent set** is a set $A \in \mathcal{I}$.
- A **base** of \mathcal{M} is a maximal independent set
- A **base of** $S \subseteq \mathcal{X}$ in \mathcal{M} is maximal independent subset of S
 - I.e. a base of the matroid after deleting \overline{S} .
- A **circuit** $S \subseteq \mathcal{X}$ is a minimal **dependent** subset of \mathcal{X}

Independent Sets, Bases, and Circuits

Consider a matroid $\mathcal{M} = (\mathcal{X}, \mathcal{I})$.

- An **independent set** is a set $A \in \mathcal{I}$.
- A **base** of \mathcal{M} is a maximal independent set
- A **base of** $S \subseteq \mathcal{X}$ in \mathcal{M} is maximal independent subset of S
 - I.e. a base of the matroid after deleting \bar{S} .
- A **circuit** $S \subseteq \mathcal{X}$ is a minimal **dependent** subset of \mathcal{X}

What are these for:

- Graphic matroid
- Linear matroid
- Uniform matroid
- Partition matroid
- Transversal matroid

Lemma

For every $S \subseteq \mathcal{X}$, all bases of S in \mathcal{M} have the same cardinality.

- Special case of $S = \mathcal{X}$: all bases of \mathcal{M} have the same cardinality.
- Should remind you of vector space dimension

Lemma

For every $S \subseteq \mathcal{X}$, all bases of S in \mathcal{M} have the same cardinality.

- Special case of $S = \mathcal{X}$: all bases of \mathcal{M} have the same cardinality.
- Should remind you of vector space dimension
- Follows directly from the exchange property.

Lemma

For every $S \subseteq \mathcal{X}$, all bases of S in \mathcal{M} have the same cardinality.

- Special case of $S = \mathcal{X}$: all bases of \mathcal{M} have the same cardinality.
- Should remind you of vector space dimension
- Follows directly from the exchange property.

The following analogue of vector space dimension is well-defined.

Rank

- The **Rank** of $S \subseteq \mathcal{X}$ in \mathcal{M} is the size of the maximal independent subsets (i.e. bases) of S .
- The rank of \mathcal{M} is the size of the bases of \mathcal{M} .
- The function $rank_{\mathcal{M}}(S) : 2^{\mathcal{X}} \rightarrow \mathbb{N}$ is called the **rank function** of \mathcal{M} .

Rank

Lemma

For every $S \subseteq \mathcal{X}$, all bases of S in \mathcal{M} have the same cardinality.

- Special case of $S = \mathcal{X}$: all bases of \mathcal{M} have the same cardinality.
- Should remind you of vector space dimension
- Follows directly from the exchange property.

The following analogue of vector space dimension is well-defined.

Rank

- The **Rank** of $S \subseteq \mathcal{X}$ in \mathcal{M} is the size of the maximal independent subsets (i.e. bases) of S .
- The rank of \mathcal{M} is the size of the bases of \mathcal{M} .
- The function $rank_{\mathcal{M}}(S) : 2^{\mathcal{X}} \rightarrow \mathbb{N}$ is called the **rank function** of \mathcal{M} .

E.g.: Graphic matroid, linear matroid, transversal matroid

Span

Given $S \subseteq \mathcal{X}$, $span(S) = \{i \in \mathcal{X} : rank(S) = rank(S \cup \{i\})\}$

- i.e. the elements which would form a circuit if added to a base of S
- e.g.: Linear matroid, graphic matroid, uniform matroid.

Span

Given $S \subseteq \mathcal{X}$, $span(S) = \{i \in \mathcal{X} : rank(S) = rank(S \cup \{i\})\}$

- i.e. the elements which would form a circuit if added to a base of S
- e.g.: Linear matroid, graphic matroid, uniform matroid.

Observation

i is selected by the greedy algorithm iff $i \notin span(\{1, \dots, i-1\})$

Operations preserving Matroidness

Given $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, consider the following operations:

- **Deletion:** For $B \subseteq \mathcal{X}$, we define $\mathcal{M} \setminus B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : S \in \mathcal{I}\}$$

- Graphic: deleting edges from the graph

Operations preserving Matroidness

Given $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, consider the following operations:

- **Deletion:** For $B \subseteq \mathcal{X}$, we define $\mathcal{M} \setminus B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : S \in \mathcal{I}\}$$

- Graphic: deleting edges from the graph
- **Disjoint union:** Given $M_1 = (\mathcal{X}_1, \mathcal{I}_1)$ and $M_2 = (\mathcal{X}_2, \mathcal{I}_2)$ with $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$, we define

$$M_1 \oplus M_2 = (\mathcal{X}_1 \cup \mathcal{X}_2, \{A_1 \cup A_2 : A_1 \in \mathcal{I}_1, A_2 \in \mathcal{I}_2\})$$

- Graphic: combining two node-disjoint graphs

Operations preserving Matroidness

Given $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, consider the following operations:

- **Deletion:** For $B \subseteq \mathcal{X}$, we define $\mathcal{M} \setminus B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : S \in \mathcal{I}\}$$

- Graphic: deleting edges from the graph
- **Disjoint union:** Given $M_1 = (\mathcal{X}_1, \mathcal{I}_1)$ and $M_2 = (\mathcal{X}_2, \mathcal{I}_2)$ with $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$, we define

$$M_1 \oplus M_2 = (\mathcal{X}_1 \cup \mathcal{X}_2, \{A_1 \cup A_2 : A_1 \in \mathcal{I}_1, A_2 \in \mathcal{I}_2\})$$

- Graphic: combining two node-disjoint graphs
- **Contraction:** Given $B \subseteq \mathcal{X}$, let $M/B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : B \cup S \in \mathcal{I}\}$$

- i.e. Think of B as always being included
- Graphic: contract the connected components of B

Operations preserving Matroidness

Given $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, consider the following operations:

- **Deletion:** For $B \subseteq \mathcal{X}$, we define $\mathcal{M} \setminus B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : S \in \mathcal{I}\}$$

- Graphic: deleting edges from the graph
- **Disjoint union:** Given $M_1 = (\mathcal{X}_1, \mathcal{I}_1)$ and $M_2 = (\mathcal{X}_2, \mathcal{I}_2)$ with $\mathcal{X}_1 \cap \mathcal{X}_2 = \emptyset$, we define

$$M_1 \oplus M_2 = (\mathcal{X}_1 \cup \mathcal{X}_2, \{A_1 \cup A_2 : A_1 \in \mathcal{I}_1, A_2 \in \mathcal{I}_2\})$$

- Graphic: combining two node-disjoint graphs
- **Contraction:** Given $B \subseteq \mathcal{X}$, let $M/B = (\mathcal{X}', \mathcal{I}')$ with $\mathcal{X}' = \mathcal{X} \setminus B$,

$$\mathcal{I}' = \{S \subseteq \mathcal{X}' : B \cup S \in \mathcal{I}\}$$

- i.e. Think of B as always being included
- Graphic: contract the connected components of B
- Others: **truncation, dual, union...**

Matroids as an Algebra of Tractable Discrete Problems

- Optimization over matroids is “easy”, in the same way that optimization over convex sets is “easy”

Matroids as an Algebra of Tractable Discrete Problems

- Optimization over matroids is “easy”, in the same way that optimization over convex sets is “easy”
- Operations preserving set convexity are analogous to operations preserving matroid structure

Matroids as an Algebra of Tractable Discrete Problems

- Optimization over matroids is “easy”, in the same way that optimization over convex sets is “easy”
- Operations preserving set convexity are analogous to operations preserving matroid structure
- Arguably, matroids and submodular functions are discrete analogues of convex sets and convex functions, respectively.
 - Less exhaustive

Outline

- 1 Matroids and The Greedy Algorithm
- 2 Basic Terminology and Properties
- 3 The Matroid Polytope**
- 4 Matroid Intersection

Viewing Matroids Polyhedrally

- As is often the case with tractable discrete problems, we can view their feasible set as a polyhedron

Viewing Matroids Polyhedrally

- As is often the case with tractable discrete problems, we can view their feasible set as a polyhedron
- For $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, the convex hull of independent sets can be written as a polytope in a natural way
 - The polytope is “solvable”, and admits a polytime separation oracle

Viewing Matroids Polyhedrally

- As is often the case with tractable discrete problems, we can view their feasible set as a polyhedron
- For $\mathcal{M} = (\mathcal{X}, \mathcal{I})$, the convex hull of independent sets can be written as a polytope in a natural way
 - The polytope is “solvable”, and admits a polytime separation oracle
- This perspective will be crucial for more advanced applications of matroids
 - Optimization of linear functions over matroid intersections
 - Optimization of submodular functions over matroids

The Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

- Assigns a variable x_i to every element i of the ground set
- Each feasible x is a fractional subset of \mathcal{X}
 - $0 \leq x_i \leq 1$ since the rank of a singleton is at most 1.

The Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

- Assigns a variable x_i to every element i of the ground set
- Each feasible x is a fractional subset of \mathcal{X}
 - $0 \leq x_i \leq 1$ since the rank of a singleton is at most 1.
- The 0-1 indicator vector x_I for independent set $I \in \mathcal{I}$ is in the above polytope

The Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

- Assigns a variable x_i to every element i of the ground set
- Each feasible x is a fractional subset of \mathcal{X}
 - $0 \leq x_i \leq 1$ since the rank of a singleton is at most 1.
- The 0-1 indicator vector x_I for independent set $I \in \mathcal{I}$ is in the above polytope
- In fact, we will show that $\mathcal{P}(\mathcal{M})$ is precisely the convex hull of independent sets \mathcal{I}

The Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

- Assigns a variable x_i to every element i of the ground set
- Each feasible x is a fractional subset of \mathcal{X}
 - $0 \leq x_i \leq 1$ since the rank of a singleton is at most 1.
- The 0-1 indicator vector x_I for independent set $I \in \mathcal{I}$ is in the above polytope
- In fact, we will show that $\mathcal{P}(\mathcal{M})$ is precisely the convex hull of independent sets \mathcal{I}
- Note: polytope has $2^{|\mathcal{X}|}$ constraints.

Integrality of the Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}(\mathcal{M}) = \text{convexhull} \{x_I : I \in \mathcal{I}\}$$

Integrality of the Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

Theorem

$$\mathcal{P}(\mathcal{M}) = \text{convexhull} \{x_I : I \in \mathcal{I}\}$$

- It is clear that $\mathcal{P}(\mathcal{M}) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}\}$

Integrality of the Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

Theorem

$$\mathcal{P}(\mathcal{M}) = \text{convexhull} \{x_I : I \in \mathcal{I}\}$$

- It is clear that $\mathcal{P}(\mathcal{M}) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}\}$
- To show that $\mathcal{P}(\mathcal{M}) \subseteq \text{convexhull} \{x_I : I \in \mathcal{I}\}$, we will show that every vertex of $\mathcal{P}(\mathcal{M})$ equals x_I for some $I \in \mathcal{I}$.

Integrality of the Matroid Polytope

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

Theorem

$$\mathcal{P}(\mathcal{M}) = \text{convexhull} \{x_I : I \in \mathcal{I}\}$$

- It is clear that $\mathcal{P}(\mathcal{M}) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}\}$
- To show that $\mathcal{P}(\mathcal{M}) \subseteq \text{convexhull} \{x_I : I \in \mathcal{I}\}$, we will show that every vertex of $\mathcal{P}(\mathcal{M})$ equals x_I for some $I \in \mathcal{I}$.
- Recall: suffices to show that every linear function $w^T x$ is maximized over $\mathcal{P}(\mathcal{M})$ at some x_I for $I \in \mathcal{I}$.

Recall: The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
- 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
- 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .

Theorem

The greedy algorithm returns the maximum weight set for every choice of weights if and only if the set system $(\mathcal{X}, \mathcal{I})$ is a matroid.

Recall: The Greedy Algorithm

- 1 $B \leftarrow \emptyset$
- 2 Sort nonnegative elements of \mathcal{X} in decreasing order of weight
 - $\{1, \dots, n\}$ with $w_1 \geq w_2 \geq \dots \geq w_n \geq 0$.
- 3 For $i = 1$ to n :
 - if $B \cup \{i\} \in \mathcal{I}$, add i to B .

Theorem

The greedy algorithm returns the maximum weight set for every choice of weights if and only if the set system $(\mathcal{X}, \mathcal{I})$ is a matroid.

- We can think of the greedy algorithm as computing the indicator vector $x^* = x_B \in \mathcal{P}(\mathcal{M})$
- We will show that x^* maximizes $w^\top x$ over $x \in \mathcal{P}(\mathcal{M})$.

Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$

- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$

- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

- Therefore, $x_i^* = \text{rank}[1 : i] - \text{rank}[1 : i-1]$

$$\sum_i w_i x_i^* = \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1])$$

Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$


- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

- Therefore, $x_i^* = \text{rank}[1 : i] - \text{rank}[1 : i-1]$

$$\sum_i w_i x_i^* = \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1])$$

- Consider an arbitrary $x \in \mathcal{P}(\mathcal{M})$

$$\sum_i w_i x_i =$$



Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$


- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

- Therefore, $x_i^* = \text{rank}[1 : i] - \text{rank}[1 : i-1]$

$$\sum_i w_i x_i^* = \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1])$$

- Consider an arbitrary $x \in \mathcal{P}(\mathcal{M})$

$$\sum_i w_i x_i = \sum_i (w_i - w_{i+1}) x(1 : i)$$



Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$


- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

- Therefore, $x_i^* = \text{rank}[1 : i] - \text{rank}[1 : i-1]$

$$\sum_i w_i x_i^* = \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1])$$

- Consider an arbitrary $x \in \mathcal{P}(\mathcal{M})$

$$\begin{aligned} \sum_i w_i x_i &= \sum_i (w_i - w_{i+1}) x(1 : i) \\ &\leq \sum_i (w_i - w_{i+1}) \text{rank}(1 : i) \end{aligned}$$



Recall: Observation

i is selected by the greedy algorithm iff $i \notin \text{span}(\{1, \dots, i-1\})$

- i.e. if $\text{rank}[1 : i] - \text{rank}[1 : i-1] = 1$.

- Therefore, $x_i^* = \text{rank}[1 : i] - \text{rank}[1 : i-1]$

$$\sum_i w_i x_i^* = \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1])$$

- Consider an arbitrary $x \in \mathcal{P}(\mathcal{M})$

$$\begin{aligned} \sum_i w_i x_i &= \sum_i (w_i - w_{i+1}) x(1 : i) \\ &\leq \sum_i (w_i - w_{i+1}) \text{rank}(1 : i) \\ &= \sum_i w_i (\text{rank}[1 : i] - \text{rank}[1 : i-1]) \end{aligned}$$



The Matroid Base Polytope

- The matroid polytope is the convex hull of independent sets
 - Graphic: convex hull of forests
- What if we wish to consider only “full-rank” sets?
 - Graphic: spanning trees

The Matroid Base Polytope

- The matroid polytope is the convex hull of independent sets
 - Graphic: convex hull of forests
- What if we wish to consider only “full-rank” sets?
 - Graphic: spanning trees

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

- The 0-1 indicator vector x_B for a base B of \mathcal{M} is in above polytope

The Matroid Base Polytope

- The matroid polytope is the convex hull of independent sets
 - Graphic: convex hull of forests
- What if we wish to consider only “full-rank” sets?
 - Graphic: spanning trees

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

- The 0-1 indicator vector x_B for a base B of \mathcal{M} is in above polytope
- In fact, we will show that $\mathcal{P}(\mathcal{M})$ is precisely the convex hull of bases of \mathcal{M}

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}_{base}(\mathcal{M}) = \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}_{base}(\mathcal{M}) = \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

- As before, one direction is obvious:

$$\mathcal{P}_{base}(\mathcal{M}) \supseteq \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}_{base}(\mathcal{M}) = \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

- As before, one direction is obvious:
 $\mathcal{P}_{base}(\mathcal{M}) \supseteq \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$
- For the other direction, take $x \in \mathcal{P}_{base}(\mathcal{M})$

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}_{base}(\mathcal{M}) = \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

- As before, one direction is obvious:
 $\mathcal{P}_{base}(\mathcal{M}) \supseteq \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$
- For the other direction, take $x \in \mathcal{P}_{base}(\mathcal{M})$
- Since $x \in \mathcal{P}(\mathcal{M})$, x is a convex combination of independent sets I_1, \dots, I_k of \mathcal{M} .

Polytope $\mathcal{P}_{base}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ \sum_{i \in \mathcal{X}} x_i &= \text{rank}(\mathcal{M}) \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

$$\mathcal{P}_{base}(\mathcal{M}) = \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$$

- As before, one direction is obvious:
 $\mathcal{P}_{base}(\mathcal{M}) \supseteq \text{convexhull} \{x_B : B \text{ is a base of } \mathcal{M}\}$
- For the other direction, take $x \in \mathcal{P}_{base}(\mathcal{M})$
- Since $x \in \mathcal{P}(\mathcal{M})$, x is a convex combination of independent sets I_1, \dots, I_k of \mathcal{M} .
- Since $\|x\|_1 = \text{rank}(\mathcal{M})$, and $\|x_{I_\ell}\|_1 \leq \text{rank}(\mathcal{M})$ for all ℓ , it must be that $\|x_{I_1}\|_1 = \|x_{I_2}\|_1 = \dots = \|x_{I_k}\|_1 = \text{rank}(\mathcal{M})$

Solvability of Matroid Polytopes

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}}(S), \quad \text{for } S \subseteq \mathcal{X}.$$
$$x_i \geq 0, \quad \text{for } i \in \mathcal{X}.$$

- When given an independence oracle for \mathcal{M} , we can maximize linear functions over $\mathcal{P}(\mathcal{M})$ in $O(n \log n)$ time
 - By integrality, same as finding max-weight independent set of \mathcal{M} .

Solvability of Matroid Polytopes

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

- When given an independence oracle for \mathcal{M} , we can maximize linear functions over $\mathcal{P}(\mathcal{M})$ in $O(n \log n)$ time
 - By integrality, same as finding max-weight independent set of \mathcal{M} .
- Therefore, by equivalence of separation and optimization, can also implement a separation oracle for $\mathcal{P}(\mathcal{M})$

Solvability of Matroid Polytopes

Polytope $\mathcal{P}(\mathcal{M})$ for $\mathcal{M} = (\mathcal{X}, \mathcal{I})$

$$\begin{aligned} \sum_{i \in S} x_i &\leq \text{rank}_{\mathcal{M}}(S), & \text{for } S \subseteq \mathcal{X}. \\ x_i &\geq 0, & \text{for } i \in \mathcal{X}. \end{aligned}$$

- When given an independence oracle for \mathcal{M} , we can maximize linear functions over $\mathcal{P}(\mathcal{M})$ in $O(n \log n)$ time
 - By integrality, same as finding max-weight independent set of \mathcal{M} .
- Therefore, by equivalence of separation and optimization, can also implement a separation oracle for $\mathcal{P}(\mathcal{M})$
- A more direct proof: reduces to **submodular function minimization**
 - $\text{rank}_{\mathcal{M}}$ is a submodular set function.

Outline

- 1 Matroids and The Greedy Algorithm
- 2 Basic Terminology and Properties
- 3 The Matroid Polytope
- 4 Matroid Intersection**

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- i.e. a set is feasible if it is independent in both matroids

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- i.e. a set is feasible if it is independent in both matroids
- In general, does not produce a matroid

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- i.e. a set is feasible if it is independent in both matroids
- In general, does not produce a matroid
- Nevertheless, it will turn out that maximizing linear functions over a matroid intersection is tractable

Matroid Intersection

- Optimization of linear functions over matroids is tractable
- Matroid operations provide an algebra for constructing new matroids from old
- We will look at one operation on matroids which does not produce a matroid, but nevertheless produces a solvable problem.

Matroid Intersection


Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- i.e. a set is feasible if it is independent in both matroids
- In general, does not produce a matroid
- Nevertheless, it will turn out that maximizing linear functions over a matroid intersection is tractable
- However, maximizing linear functions over the intersection of 3 or more matroids is NP-hard

Examples

Bipartite Matching


Given a bipartite graph G , a set of edges F is a bipartite matching if and only if each node is incident on at most one edge in F .



Examples


Bipartite Matching

Given a bipartite graph G , a set of edges F is a bipartite matching if and only if each node is incident on at most one edge in F .



Arborescence


Given a directed graph G , a set of edges is an r -arborescence if it is a tree directed away from the root r .



Examples


Bipartite Matching

Given a bipartite graph G , a set of edges F is a bipartite matching if and only if each node is incident on at most one edge in F .



Arborescence

Given a directed graph G , a set of edges is an r -arborescence if it is a tree directed away from the root r .



- Others: colorful spanning trees, ...

The Matroid Intersection Polytope

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- Optimizing a modular function over $\mathcal{M}_1 \cap \mathcal{M}_2$ is equivalent to optimizing a linear function over $\text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$.
- As it turns out, this is a solvable polytope.

Theorem

$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) = \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

The Matroid Intersection Polytope

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- Optimizing a modular function over $\mathcal{M}_1 \cap \mathcal{M}_2$ is equivalent to optimizing a linear function over $\text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$.
- As it turns out, this is a solvable polytope.

Theorem

$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) = \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

- One direction is obvious:
$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

The Matroid Intersection Polytope

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- Optimizing a modular function over $\mathcal{M}_1 \cap \mathcal{M}_2$ is equivalent to optimizing a linear function over $\text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$.
- As it turns out, this is a solvable polytope.

Theorem

$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) = \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

- One direction is obvious:
 $\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$
- The other direction is not so obvious

The Matroid Intersection Polytope

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- Optimizing a modular function over $\mathcal{M}_1 \cap \mathcal{M}_2$ is equivalent to optimizing a linear function over $\text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$.
- As it turns out, this is a solvable polytope.

Theorem

$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) = \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

- One direction is obvious:
 $\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$
- The other direction is not so obvious
 - It is conceivable that $\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2)$ has fractional vertices

The Matroid Intersection Polytope

Matroid Intersection

Given matroids $\mathcal{M}_1 = (\mathcal{X}, \mathcal{I}_1)$ and $\mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_2)$ on the same ground set, we define the set system $\mathcal{M}_1 \cap \mathcal{M}_2 = (\mathcal{X}, \mathcal{I}_1 \cap \mathcal{I}_2)$.

- Optimizing a modular function over $\mathcal{M}_1 \cap \mathcal{M}_2$ is equivalent to optimizing a linear function over $\text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$.
- As it turns out, this is a solvable polytope.

Theorem

$$\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) = \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$$

- One direction is obvious:
 $\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2) \supseteq \text{convexhull} \{x_I : I \in \mathcal{I}_1 \cap \mathcal{I}_2\}$
- The other direction is not so obvious
 - It is conceivable that $\mathcal{P}(\mathcal{M}_1) \cap \mathcal{P}(\mathcal{M}_2)$ has fractional vertices
- Nevertheless, it is true but hard to prove, so we will skip it.

Optimization over Matroid Intersection $\mathcal{M}_1 \cap \mathcal{M}_2$

$$\begin{array}{ll} \text{maximize} & \sum_{i \in \mathcal{X}} w_i x_i \\ \text{subject to} & \\ & \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_1}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_2}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & x_i \geq 0, \quad \text{for } i \in \mathcal{X}. \end{array}$$

Optimization over Matroid Intersections

Optimization over Matroid Intersection $\mathcal{M}_1 \cap \mathcal{M}_2$

$$\begin{aligned} & \text{maximize} && \sum_{i \in \mathcal{X}} w_i x_i \\ & \text{subject to} && \\ & && \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_1}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & && \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_2}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & && x_i \geq 0, \quad \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

Given independence oracles to both matroids \mathcal{M}_1 and \mathcal{M}_2 , there is an algorithm for finding the maximum weight set in $\mathcal{M}_1 \cap \mathcal{M}_2$ which runs in $\text{poly}(n)$ time.

Optimization over Matroid Intersections

Optimization over Matroid Intersection $\mathcal{M}_1 \cap \mathcal{M}_2$

$$\begin{aligned} & \text{maximize} && \sum_{i \in \mathcal{X}} w_i x_i \\ & \text{subject to} && \\ & && \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_1}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & && \sum_{i \in S} x_i \leq \text{rank}_{\mathcal{M}_2}(S), \quad \text{for } S \subseteq \mathcal{X}. \\ & && x_i \geq 0, \quad \text{for } i \in \mathcal{X}. \end{aligned}$$

Theorem

Given independence oracles to both matroids \mathcal{M}_1 and \mathcal{M}_2 , there is an algorithm for finding the maximum weight set in $\mathcal{M}_1 \cap \mathcal{M}_2$ which runs in $\text{poly}(n)$ time.

Proof: Using equivalence of separation and optimization, and the fact that all coefficients in the LP have $\text{poly}(n)$ bits.

NP-hardness of 3-way Matroid Intersection

By a reduction from **Hamiltonian Path** in directed graphs