


CS675: Convex and Combinatorial Optimization
Spring 2022
Geometric Duality of Convex Sets and Functions

Instructor: Shaddin Dughmi

Outline

- 1 Convexity and Duality
- 2 Duality of Convex Sets
- 3 Duality of Convex Functions


Duality Correspondences



There are two equivalent ways to represent a convex set

- The family of points in the set (standard representation)
- The set of halfspaces containing the set (“dual” representation)

Duality Correspondances




There are two equivalent ways to represent a convex set

- The family of points in the set (standard representation)
- The set of halfspaces containing the set (“dual” representation)

This equivalence between the two representations gives rise to a variety of “duality” relationships among convex sets, cones, and functions.

Duality Correspondances




There are two equivalent ways to represent a convex set

- The family of points in the set (standard representation)
- The set of halfspaces containing the set (“dual” representation)

This equivalence between the two representations gives rise to a variety of “duality” relationships among convex sets, cones, and functions.


Definition

“**Duality**” is a woefully overloaded mathematical term for a relation that groups elements of a set into “dual” pairs.



Theorem

A closed convex set S is the intersection of all closed halfspaces \mathcal{H} containing it.




Theorem

A closed convex set S is the intersection of all closed halfspaces \mathcal{H} containing it.


Proof

- Clearly, $S \subseteq \bigcap_{H \in \mathcal{H}} H$
- To prove equality, consider $x \notin S$
- By the separating hyperplane theorem, there is a hyperplane separating S from x
- Therefore there is $H \in \mathcal{H}$ with $x \notin H$, hence $x \notin \bigcap_{H \in \mathcal{H}} H$



Theorem

A closed convex cone K is the intersection of all closed homogeneous halfspaces \mathcal{H} containing it.




Theorem

A closed convex cone K is the intersection of all closed homogeneous halfspaces \mathcal{H} containing it.


Proof

- For every non-homogeneous halfspace $a^\top x \leq b$ containing K , the smaller homogeneous halfspace $a^\top x \leq 0$ contains K as well.
- Therefore, can discard non-homogeneous halfspaces without changing the intersection



Theorem

A convex function is the point-wise supremum of all affine functions under-estimating it everywhere.



Theorem

A convex function is the point-wise supremum of all affine functions under-estimating it everywhere.



Proof

- **epi** f convex, therefore is the intersection of family of halfspaces \mathcal{H}
- Each $h \in \mathcal{H}$ can be written as $a^\top x - t \leq b$, for some $a \in \mathbb{R}^n$ and $b \in \mathbb{R}$. (Why?)
 - Constrains $(x, t) \in \mathbf{epi} f$ to $a^\top x - b \leq t$
- $f(x)$ is the lowest t s.t. $(x, t) \in \mathbf{epi} f$
- Therefore, $f(x)$ is the point-wise maximum of $a^\top x - b$ over all halfspaces $h(a, b) \in \mathcal{H}$.

Outline

- 1 Convexity and Duality
- 2 Duality of Convex Sets**
- 3 Duality of Convex Functions

Polar Duality of Convex Sets



One way of representing all the halfspaces containing a convex set.

Polar

Let $S \subseteq \mathbb{R}^n$ be a closed convex set containing the origin. The **polar** of S is defined as follows:

$$S^\circ = \{y : y^\top x \leq 1 \text{ for all } x \in S\}$$

Note

- Every halfspace $a^\top x \leq b$ with $b \neq 0$ can be written as a “normalized” inequality $y^\top x \leq 1$, by dividing by b .
- S° can be thought of as the normalized representations of halfspaces containing S .

$$S^\circ = \{y : y^\top x \leq 1 \text{ for all } x \in S\}$$

Properties of the Polar

- 1 $S^{\circ\circ} = S$
- 2 S° is a closed convex set containing the origin
- 3 When 0 is in the interior of S , then S° is bounded.

$$S^\circ = \{y : y^\top x \leq 1 \text{ for all } x \in S\}$$

Properties of the Polar

- 1 $S^{\circ\circ} = S$
- 2 S° is a closed convex set containing the origin
- 3 When 0 is in the interior of S , then S° is bounded.

- 2 Follows from representation as intersection of halfspaces
- 3 S contains an ϵ -ball centered at the origin, so S° is contained in the $\frac{1}{\epsilon}$ ball centered at the origin.
 - Take $y \in S^\circ$
 - $x := \epsilon \frac{y}{\|y\|_2} \in S$
 - $1 \geq y^\top x = \epsilon \|y\|_2$, so $\|y\|_2 \leq \frac{1}{\epsilon}$

$$S^\circ = \{y : y^\top x \leq 1 \text{ for all } x \in S\}$$

Properties of the Polar

- 1 $S^{\circ\circ} = S$
- 2 S° is a closed convex set containing the origin
- 3 When 0 is in the interior of S , then S° is bounded.

$$S^{\circ\circ} = \{x : x^\top y \leq 1 \text{ for all } y \in S^\circ\}$$

- 1
 - $S \subseteq S^{\circ\circ}$ is easy: $\hat{x} \in S \implies \forall y \in S^\circ \hat{x}^\top y \leq 1 \implies \hat{x} \in S^{\circ\circ}$
 - Take $\hat{x} \notin S$, by SSHT and $0 \in S$, there is a halfspace $z^\top x \leq 1$ containing S but not \hat{x} (i.e. $z^\top \hat{x} > 1$)
 - $z \in S^\circ$, therefore $\hat{x} \notin S^{\circ\circ}$

$$S^\circ = \{y : y^T x \leq 1 \text{ for all } x \in S\}$$

Properties of the Polar

- 1 $S^{\circ\circ} = S$
- 2 S° is a closed convex set containing the origin
- 3 When 0 is in the interior of S , then S° is bounded.

Note

When S is non-convex, $S^\circ = (\text{convexhull}(S))^\circ$, and $S^{\circ\circ} = \text{convexhull}(S)$.

Examples




The unit sphere for different metrics: $\|x\|_{l_p} = 1$ in \mathbb{R}^2 .

Norm Balls

- The polar of the Euclidean unit ball is itself (we say it is **self-dual**)
- The polar of the 1-norm ball is the ∞ -norm ball
- More generally, the p -norm ball is dual to the q -norm ball, where

$$\frac{1}{p} + \frac{1}{q} = 1$$




Polytopes

Given a polytope P represented as $Ax \preceq \vec{1}$, the polar P° is the convex hull of the rows of A .

- Facets of P correspond to vertices of P° .
- Dually, vertices of P correspond to facets of P° .

Polar Duality of Convex Cones



Polar duality takes a simplified form when applied to cones

Polar


The polar of a closed convex cone K is given by

$$K^\circ = \{y : y^\top x \leq 0 \text{ for all } x \in K\}$$

Note

- $\forall x \in K \ y^\top x \leq 1 \iff \forall x \in K \ y^\top x \leq 0$
- K° represents all homogeneous halfspaces containing K .

Polar Duality of Convex Cones



Polar duality takes a simplified form when applied to cones

Polar

The polar of a closed convex cone K is given by

$$K^\circ = \{y : y^\top x \leq 0 \text{ for all } x \in K\}$$

Dual Cone

By convention, $K^* = -K^\circ$ is referred to as the **dual cone** of K .

$$K^* = \{y : y^\top x \geq 0 \text{ for all } x \in K\}$$

$$K^\circ = \{y : y^\top x \leq 0 \text{ for all } x \in K\}$$

Properties of the Polar Cone

- 1 $K^{\circ\circ} = K$
- 2 K° is a closed convex cone

$$K^\circ = \{y : y^\top x \leq 0 \text{ for all } x \in K\}$$

Properties of the Polar Cone


- 1 $K^{\circ\circ} = K$
 - 2 K° is a closed convex cone
-
- 1 Same as before
 - 2 Intersection of homogeneous halfspaces

Examples

- The polar of a subspace is its orthogonal complement
- The polar cone of the nonnegative orthant is the nonpositive orthant
 - Self-dual
- The polar of a polyhedral cone $Ax \preceq 0$ is the conic hull of the rows of A
- The polar of the cone of positive semi-definite matrices is the cone of negative semi-definite matrices
 - Self-dual

Recall: Farkas' Lemma

Let K be a **closed convex cone** and let $w \notin K$. There is $z \in \mathbb{R}^n$ such that $z^\top x \leq 0$ for all $x \in K$, and $z^\top w > 0$.




Equivalently: there is $z \in K^\circ$ with $z^\top w > 0$.

Outline

- 1 Convexity and Duality
- 2 Duality of Convex Sets
- 3 Duality of Convex Functions**

Conjugation of Convex Functions




Conjugate

For a function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$, the **conjugate** of f is

$$f^*(y) = \sup_x (y^\top x - f(x))$$

Note


- $f^*(y)$ is the minimal value of β such that the affine function $y^\top x - \beta$ underestimates $f(x)$ everywhere.
- Equivalently, the distance we need to lower the hyperplane $y^\top x - t = 0$ in order to get a supporting hyperplane to $\text{epi } f$.
- $y^\top x - t = f^*(y)$ are the supporting hyperplanes of $\text{epi } f$



$$f^*(y) = \sup_x (y^\top x - f(x))$$

Properties of the Conjugate

- 1 $f^{**} = f$ when f is convex
- 2 f^* is a convex function
- 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)




$$f^*(y) = \sup_x (y^T x - f(x))$$

Properties of the Conjugate

- 1 $f^{**} = f$ when f is convex
- 2 f^* is a convex function
- 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)

- 2 Supremum of affine functions of y
- 3 By definition of $f^*(y)$




$$f^*(y) = \sup_x (y^\top x - f(x))$$

Properties of the Conjugate

- 1 $f^{**} = f$ when f is convex
- 2 f^* is a convex function
- 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)


- 1 • $f^{**}(x) = \max_y y^\top x - f^*(y)$ by definition



$$f^*(y) = \sup_x (y^\top x - f(x))$$

Properties of the Conjugate


- 1 $f^{**} = f$ when f is convex
 - 2 f^* is a convex function
 - 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)
- 1
- $f^{**}(x) = \max_y y^\top x - f^*(y)$ by definition
 - For fixed y , $f^*(y)$ is minimal β such that $y^\top x - \beta$ underestimates f .



$$f^*(y) = \sup_x (y^T x - f(x))$$

Properties of the Conjugate

- 1 $f^{**} = f$ when f is convex
- 2 f^* is a convex function
- 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)
 - 1
 - $f^{**}(x) = \max_y y^T x - f^*(y)$ by definition
 - For fixed y , $f^*(y)$ is minimal β such that $y^T x - \beta$ underestimates f .
 - Therefore $f^{**}(x)$ is the maximum, over all y , of affine underestimates $y^T x - \beta$ of f



$$f^*(y) = \sup_x (y^\top x - f(x))$$

Properties of the Conjugate

- 1 $f^{**} = f$ when f is convex
- 2 f^* is a convex function
- 3 $xy \leq f(x) + f^*(y)$ for all $x, y \in \mathbb{R}^n$ (Fenchel's Inequality)

- 1
 - $f^{**}(x) = \max_y y^\top x - f^*(y)$ by definition
 - For fixed y , $f^*(y)$ is minimal β such that $y^\top x - \beta$ underestimates f .
 - Therefore $f^{**}(x)$ is the maximum, over all y , of affine underestimates $y^\top x - \beta$ of f
 - By our earlier characterization, this is equal to f when f is convex.

Examples

- Affine function: $f(x) = ax + b$. Conjugate has $f^*(a) = -b$, and ∞ elsewhere
- $f(x) = x^2/2$ is self-conjugate
- Exponential: $f(x) = e^x$. Conjugate has $f^*(y) = y \log y - y$ for $y \in \mathbb{R}_+$, and ∞ elsewhere.
- Convex Quadratic: $f(x) = \frac{1}{2}x^\top Qx$ with Q positive definite. Conjugate is $f^*(y) = \frac{1}{2}y^\top Q^{-1}y$
- Log-sum-exp: $f(x) = \log(\sum_i e^{x_i})$. Conjugate has $f^*(y) = \sum_i y_i \log y_i$ for $y \succeq 0$ and $1^\top y = 1$, ∞ otherwise.